

On the signed graphs with two distinct eigenvalues

F. Ramezani ¹

Department of Mathematics,

K.N.Toosi University of Technology, Tehran, Iran

P.O. Box 16315-1618

Abstract. We consider signed graphs, i.e, graphs with positive or negative signs on their edges. We construct some families of bipartite signed graphs with only two distinct eigenvalues. This leads to constructing infinite families of regular Ramanujan graphs.

Keywords: Signed graphs, two distinct eigenvalues, Ramanujan graphs.

Mathematics Subject Classification: 05C50.

1 Introduction

We consider only simple graphs, i.e., graphs with out loops and multiple edges. The vertex set and edge set of the graph G will be denoted by $V(G)$ and $E(G)$, respectively. If there is no doubt about G we simply write V and E . A *signature* on a graph G is a function $s : E \rightarrow \{1, -1\}$. A graph G provided with a signature s is called a *signed graph*, and will be denoted by $G(s)$. We call the graph G the *ground graph* of the signed graph $G(s)$. The *adjacency matrix*, A^s of the signed graph $G(s)$ on the vertex set $V = \{v_1, v_2, \dots, v_n\}$, is

¹Email: ramezani@kntu.ac.ir.

an $n \times n$ matrix whose entries are

$$A^s(i, j) = \begin{cases} 1, & \text{if } v_i \text{ is adjacent to } v_j \text{ and } s(\{v_i, v_j\}) = 1; \\ -1, & \text{if } v_i \text{ is adjacent to } v_j \text{ and } s(\{v_i, v_j\}) = -1; \\ 0, & \text{otherwise.} \end{cases}$$

The $n \times n$ matrix A , whose entries are the absolute values of the entries of A^s , is called the *ordinary adjacency matrix* of the graph G and will be denote by $A(G) = |A^s|$. Note that any symmetric $(0, \pm 1)$ -matrix with zero diagonal is corresponding to a signature of the graph whose adjacency matrix is $|C|$. Then we some times use the symmetric matrix C rather than the signature which is in it's correspondence. The *spectrum* of a signed graph is the eigenvalues of it's adjacency matrix. By O_n , J_n and I_n we mean the all zero matrix, all one matrix, and identity matrix of size n , respectively. If there is no doubt about size of the matrices we simply write O, J, I . For a signed matrix A of order n , by A^* we mean the matrix $\begin{pmatrix} O_n & A \\ A' & O_n \end{pmatrix}$, where A' is the transpose of the matrix A . An $n \times n$ matrix C is called *orthogonal signed matrix* if the entries of C belongs to the set $\{0, 1, -1\}$ and $CC' = C'C = \alpha I_n$, where α is a positive integer. A signed graph is called *orthogonal* if it's adjacency matrix is an orthogonal signed matrix. Recently some problems on the spectrum of signed adjacency matrices have attracted many studies. In [2], the authors have considered the lollipop graph and proved it's signed graphs are determined by their spectrum. Energy of signed matrices have been considered in [5]. The spectrum of signed graphs can be used to find the spectrum of 2-lifts of graphs, which leads to some results on the existence of infinite families of regular Ramanujan graphs with a fixed degree (cf. [7]). It is known that the only graphs with two distinct eigenvalue of the ordinary adjacency matrix are the complete graphs. In this article we consider bipartite graphs and find some signatures of them which leads to signed graphs with only two distinct eigenvalues. In [8] the authors have proved the existence of infinite families of regular bipartite Ramanujan graphs of every degree bigger than 2. Here using the Hadamard matrices we construct some of regular bipartite Ramanujan graphs. During the preparation of this paper we encounter interesting results from other combinatorial areas such as regular two-graphs, Hadamard matrices, Conference matrices and Orthogonal codes.

2 preliminaries

In this section we bring some basic tools and results which we will need to construct the signed graphs with only two distinct eigenvalues. We must look for the symmetric matrices with entries $0, \pm 1$ which holds in an integral monic polynomial of degree 2. In [10] the complete signed graphs with two distinct eigenvalues are characterized. There, it is proved that a two-graph is regular if and only if it's (signed) adjacency matrix has just two distinct eigenvalues.

In a signed graph $G(s)$ *resigning* around the vertex $v \in V(G)$ is multiplying the signs of all the edges incident to v by -1 . Two signed graphs $G(s)$ and $G(s')$ are called *equivalent* if one is obtained from the other by a sequence of resigning around prescribed vertices. Otherwise we call them *distinct*. Most properties of signed graphs are preserved under resigning around a vertex. In the following proposition from [9] the number of distinct signed graphs on the ground graph G is enumerated.

Proposition 1 [9] If G has m edges, n vertices and c components, then there are $2^{(m-n+c)}$ distinct signed graphs on G .

In the following lemmas we mention the basic properties of possible signed graphs with just two distinct eigenvalues.

Lemma 1 Let s be a signature on the graph G , such that the signed graph $G(s)$ has just two distinct eigenvalues, then the graph G should be regular.

Proof. The diagonal entries of the matrix $(A^s)^2$ are the vertex degrees of G . On the other hand if λ, μ are the distinct eigenvalues of $G(s)$, then $(A^s)^2 - (\lambda + \mu)A^s + \lambda\mu I = 0$. The diagonal entries of the matrix A^s are zero hence the diagonal entries of $(A^s)^2$ (therefore the vertex degrees of G) should all be equal to $-\lambda\mu$, this implies that G is regular. $\square$

Lemma 2 Let G be a bipartite graph on $2n$ vertices, and s be a signature of it, then the signed graph $G(s)$ has only two distinct eigenvalues if and only if it's adjacency matrix is of the form $\begin{pmatrix} 0 & C \\ C' & 0 \end{pmatrix}$, where C is an orthogonal signed matrix of size n .

Proof. If G is a bipartite graph on $2n$ vertices, then the adjacency matrix of $G(s)$ is of the block form $\begin{pmatrix} 0 & C \\ C' & 0 \end{pmatrix}$, on the other hand we have $(A^s)^2 + \alpha A^s + \beta I_{2n} = 0$, for some integers α, β . It implies

$$\begin{pmatrix} 0 & C \\ C' & 0 \end{pmatrix}^2 + \alpha \begin{pmatrix} 0 & C \\ C' & 0 \end{pmatrix} + \beta I_{2n} = 0.$$

Therefore,

$$\begin{pmatrix} CC' & 0 \\ 0 & C'C \end{pmatrix} + \alpha \begin{pmatrix} 0 & C \\ C' & 0 \end{pmatrix} + \beta I_{2n} = 0.$$

The above equality yields

$$CC' = C'C = \beta I_n, \quad \alpha C = 0.$$

Hence $\alpha = 0$ and C and A^s are orthogonal signed matrices. On the other hand suppose that C is an orthogonal signed matrix of size n , with $CC' = C'C = \alpha I$. Then for the symmetric signed matrix $A^s = C^*$, we have $(A^s)^2 = \alpha I_{2n}$ and hence the corresponding signed graph has just two distinct eigenvalues $\pm\sqrt{\alpha}$. $\square$

The above lemma implies that any bipartite signed graph with just two distinct eigenvalues is in correspondence with an orthogonal signed matrix C . Therefore from now on we will refer to orthogonal signed matrices instead of bipartite signed graphs with just two distinct eigenvalues. We will denote the corresponding orthogonal signed matrix of the signed graph $G(s)$ by $C_s(G)$.

Corollary 1 If G is a complete bipartite graph with $2n$ vertices, and $G(s)$ has only two distinct eigenvalues, then the matrix $C_s(G)$ is an Hadamard matrix of size n .

Corollary 2 If G is a bipartite $(n - 1)$ -regular graph on $2n$ vertices, and $G(s)$ has only two distinct eigenvalues, then the matrix $C_s(G)$ is a Conference matrix of size n .

3 Constructing bipartite signed graphs with two distinct eigenvalues

In this section we first recall some well known methods of constructing the Hadamard matrices, then apply them to find orthogonal signed matrices and hence bipartite signed graphs with just two distinct eigenvalues. We bring the following lemma with out proof.

Lemma 3 Let A and B are orthogonal matrices, then the matrix $A \otimes B$, which is the Kronecker product of A, B , is orthogonal.

The following corollaries are immediate consequences of the above lemma.

Corollary 3 If there exist bipartite signed graphs on respectively $2n, 2m$ vertices, with only two distinct eigenvalues, then there exists a bipartite signed graph on $4mn$ vertices, which has only two distinct eigenvalues.

Corollary 4 For any positive integer n , there is an orthogonal signed matrix of size $2n$, and hence a bipartite signed graph on $4n$ vertices and only two distinct eigenvalues

Proof. Consider the Kronecker product of the matrix I_n and H_2 , where H_2 is the Hadamard matrix of size 2, i.e

$$H_2 = \begin{pmatrix} 1 & 1 \\ 1 & -1 \end{pmatrix}.$$

□

Lemma 4 Let C be a symmetric $n \times n$ orthogonal signed matrix with zero entries on the diagonal. Then the matrix defined bellow is a symmetric orthogonal signed matrix,

$$B = \begin{pmatrix} C + I & C - I \\ C - I & -C - I \end{pmatrix}.$$

Proof. We have

$$B' = \begin{pmatrix} C' + I & C' - I \\ C' - I & -C' - I \end{pmatrix},$$

since C is symmetric so is B . Suppose that $CC' = C'C = \alpha I$, we have

$$\begin{aligned} BB' &= \begin{pmatrix} (C+I)(C'+I) + (C-I)(C'-I) & (C+I)(C'-I) + (C-I)(-C'-I) \\ (C-I)(C'+I) + (-C-I)(C'-I) & (C-I)(C'-I) + (-C-I)(-C'-I) \end{pmatrix}, \\ &= \begin{pmatrix} CC' + 2I & 0 \\ 0 & CC' + 2I \end{pmatrix} = \begin{pmatrix} (\alpha+2)I & 0 \\ 0 & (\alpha+2)I \end{pmatrix}, \end{aligned}$$

so the assertion follows. $\square$

Lemma 5 Let C be an antisymmetric orthogonal matrix with zero entries on the diagonal. Then the matrix $C + I$ is an orthogonal matrix.

Proof. We have

$$(C + I)(C' + I) = CC' + C + C' + I,$$

since C is an antisymmetric orthogonal signed matrix, the assertion follows. $\square$

In the following theorem, we establish a method for constructing orthogonal matrices, based on the Williamson method (cf. [6]).

Theorem 1 Let the matrices $A_i, 1 \leq i \leq 4$, be symmetric of order n , with entries $0, \pm 1$, having constant number of zeros in each row and assume that they are mutually commuting with each other. Consider the matrix H defined bellow,

$$H = \begin{pmatrix} A_1 & A_2 & A_3 & A_4 \\ -A_2 & A_1 & -A_4 & A_3 \\ -A_3 & A_4 & A_1 & -A_2 \\ -A_4 & -A_3 & A_2 & A_1 \end{pmatrix},$$

then H is an orthogonal matrix if and only if

$$\sum_{i=1}^4 A_i^2 = \left(\sum_{i=1}^4 k_i \right) I.$$

Where k_i is the number of non-zero entries in each row of A_i , for $i = 1, \dots, 4$.

Proof. The assertion follows by simply calculating HH' . $\square$

Proposition 2 Let C be a symmetric orthogonal signed matrix of order n . Then the matrix H , in the previous theorem, obtained by substituting any of the followings, is an orthogonal signed matrix.

- $A_1 = A_2 = A_3 = A_4 = C$, or
- $A_1 = A_2 = C$, $A_3 = C - I$ and $A_4 = C + I$, or
- $A_1 = A_2 = C + I$, $A_3 = A_4 = C - I$,

providing the matrix C has zero diagonal in the second and last cases.

Proposition 3 If the matrices A_i in Theorem 1 all be equal to a given orthogonal matrix C , the matrix H will be orthogonal.

Proof. In general the matrices A_i , defined in Theorem 1 don't need to be symmetric. By calculation we see that the only requirement for the orthogonality of H is that the matrices $A_i A_j'$ for $i, j = 1, \dots, 4$ to be symmetric. Since we have $CC' = \alpha I$, then the assertion follows. $\square$

One may consider the matrix C in the previous propositions to be a symmetric conference matrix. An infinite family of symmetric conference matrices exist, for instance with Paley method we may construct symmetric conference matrices of sizes $q + 1$, where q is a prime power and $q \equiv 1 \pmod{4}$. (cf. [6], pp. 175–176)

4 Some Ramanujan graphs

Bilu and Linial conjectured that every regular Ramanujan graph of degree d has a signature s on the edge set with the property $\lambda_1(A^s) \leq 2\sqrt{d-1}$. We will call the mentioned signature a good signature. In this section for some families of known Ramanujan graphs, we verify the conjecture of Bilu and Linial and introduce new families of Ramanujan graphs. To state our main result we need some lemmas.

Lemma 6 If G is a k -regular graph on n vertices and $\frac{1}{4}(k-1)^2 + k + 2 \leq n$ then the complement of G , G^c is a Ramanujan graph.

Proof. The graph G^c is a $(n-k-1)$ -regular graph, hence we need to prove that

$$\lambda_2(G) \leq 2\sqrt{n-k-2}.$$

It is known that the eigenvalues of the complement of G is in decreasing order as the following list,

$$n-k-1, -1-\lambda_n(G), -1-\lambda_{n-1}(G), \dots, -1-\lambda_2(G).$$

By Perron-Frobenius Theorem we know that $\lambda_n(G) \geq -k$ then $\lambda_2(G^c) \leq k-1$, and the assertion follows by the following inequalities

$$\lambda_2(G^c) \leq k-1 \Leftrightarrow \frac{1}{4}\lambda_2^2(G^c) \leq \frac{1}{4}(k-1)^2 \leq n-(k+2),$$

hence

$$\lambda_2(G^c) \leq 2\sqrt{n-k-2}$$

thus G^c is a Ramanujan graph. $\square$

Proposition 4 Let C be a symmetric orthogonal signed matrix of order n , with $CC' = \alpha I$, $\alpha \geq 1$ and $k = n-1-\alpha$, holds in the inequality $\frac{1}{4}(k-1)^2 + k + 2 \leq n$ then the ground graph G corresponding to the matrix C is a Ramanujan graph and C is a good signature of G .

Proof. The graph G is α -regular, on the other hand C is a symmetric orthogonal signed matrix hence $\lambda_1(C) = \sqrt{\alpha} \leq 2\sqrt{\alpha-1}$. The complement of G is a regular graph of degree $k = n-1-\alpha$. By the assumption and Lemma 6 the assertion follows. $\square$

In the case that the orthogonal signed matrix C is not symmetric we can use it to construct bipartite Ramanujan graphs. But before that we need some preliminaries. For a bipartite graph G with bipartition X, Y of vertices, the *bipartite complement* of G which is denoted by G_b^c is a bipartite graph on the vertex set $V = X \cup Y$, where a vertex $x \in X$ is adjacent to a vertex $y \in Y$ if and only if x was not adjacent to y in G .

Lemma 7 If G is a k -regular bipartite Ramanujan graph on the disjoint sets X, Y of size n , and $k \leq \frac{n}{2}$ then the bipartite complement of G is also a Ramanujan graph.

Proof. We have $A(G) + A(G_b^c) = A(K_{n,n})$. We claim that $A(G)$ and $A(K_{n,n})$ commute and hence the eigenvalues of G_b^c follows

$$\lambda_i(G_b^c) = \lambda_i(K_{n,n}) - \lambda_i(G) = \begin{cases} n - k & i = 1, \\ \lambda_i(G) & 1 < i < n, \\ k - n & i = n. \end{cases}$$

By the assumption $\lambda_2(G) \leq 2\sqrt{k-1}$, and $k \leq n - k$ hence

$$\lambda_2(G_b^c) = \lambda_2(G) \leq 2\sqrt{k-1} \leq 2\sqrt{n-k-1},$$

therefore G_b^c is a Ramanujan graph. □

Lemma 8 The bipartite complement of the graph kC_4 , ($k \geq 2$) that is the k disjoint copy of C_4 , is a Ramanujan graph.

Proof. As discussed in the proof of the above lemma, the eigenvalues of $(kC_4)_b^c$ can be calculated by the eigenvalues of the graph $K_{2k,2k}$ and the graph kC_4 . Thus the spectrum of $(kC_4)_b^c$ is $\pm(2k-2), (\pm 2)^{k-1}, 0^{2k}$, hence the assertion follows. □

Lemma 9 If C is an orthogonal signed matrix of size n and the ground graph G corresponding to C^* is a Ramanujan graph, then C^* is a good signature of G .

For a signature s on the edges of the graph G the 2-lift $\tilde{G}_s$ of the graph G is a graph on the vertex set $V(G) \times \{1, 2\}$, where there is an edge between (u, i) and (v, j) if and only if

$$\{u, v\} \in E(G) \text{ and } \begin{cases} i = j & \text{if } s(uv) = 1, \\ i = \bar{j} & \text{otherwise.} \end{cases},$$

where $\bar{1} = 2$ and $\bar{2} = 1$. In Figure 1, we illustrate the definition. Note that in the figure the negative edge is denoted by dot line. In the graph $\tilde{G}_s$, the vertices of $G(s)$ are duplicated

and the positive edges are replaced by two parallel lines and the negative edge is replaced by two crossed edges.




Figure1. 2-lift of G corresponding to the specific signature

let $P_2 = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}$. The spectrum of the graph $\tilde{G}_s$ is determined in [7].

Lemma 10 [7] The spectrum of $A(\tilde{G}_s)$ is the multi set union of spectrum of A and A^s .

By the above Lemma the authors of the paper [8] conclude that for any Ramanujan graph G and a good signature s of it, the graph $\tilde{G}_s$ is a Ramanujan graph. As a final result, in the following table we list some Ramanujan graphs and a good signature of them. Suppose that H_n is a Hadamard matrix and Co_n is a conference matrix of order n .

G	a good signature A^s	eigenvalues of $\tilde{G}_s$
$K_{n,n}$	H_n^*	$\pm n, \pm \sqrt{n}^n, 0^{2n-2}$
$K_{n,n} \setminus M$	Co_n^*	$\pm(n-1), \pm \sqrt{n-1}^n, \pm 1^{n-1}$
$(nC_4)_b^c$	$\begin{pmatrix} Co_n & Co_n \\ -Co_n & Co_n \end{pmatrix}^*$	$\pm(2n-2), (\pm 2)^{n-1}, 0^{2n}$

Acknowledgment We would like to thank B. Tayfeh-Rezaie for attracting our opinion to the problem.

References

- [1] B.D. ACHARYA, Spectral criterion for cycle balance in networks, *J. Graph Theory* **4**(1) (1980) 1–11.
- [2] F. BELARDO AND P. PETECKI, Spectral characterizations of signed lollipop graphs *Linear Algebra and its Applications*, **480** (2015) 144–167.
- [3] EDWIN R. VAN DAM , Regular Graphs With Four Eigenvalues *Linear Algebra and its Applications*, **226-228** (1995) 139–162.
- [4] M. DOOB AND W. H. HAEMERS, The complement of the path is determined by its spectrum, *Linear Algebra and its Applications*, **356** (2002) 57-65.
- [5] K.A. GERMINA, SHAHUL HAMEED K AND THOMAS ZASLAVSKY, On products and line graphs of signed graphs, their eigenvalues and energy. *Linear Algebra and its Applications*, **435** (2011) 2432-2450.
- [6] J.H. VAN LINT AND R.M. WILLSON, *A Course in Combinatorics*, Cambridge university press, 1992.
- [7] S. LIU, N. PEYERIMHOFF, AND A. VDOVINA, *Signatures, lifts, and eigenvalues of graphs*, arXiv:1412.6841 [math.CO].
- [8] ADAM W. MARCUS, DANIEL A. SPIELMAN AND NIKHIL SRIVASTAVA *Interlacing Families I: Bipartite Ramanujan Graphs of All Degrees*, arXiv:1304.4132 [math.CO].
- [9] REZA NASERASR, EDITA ROLLOVA, AND ERIC SOPENAC, Homomorphisms of signed graphs, *J of Graph Theory* DOI 10.1002/jgt
- [10] D. E. TAYLOR, Regular 2-graphs, *Proc. London Math. Soc.* **35** (1977) 257–274.